

Holotypes of fossil species

| <u>Taxon</u> | <u>? coll.</u> | <u>Specimen</u> | <u>Stratigraphy</u> | <u>Location</u> | <u>Georeference Bibliography</u> | <u>3D-model</u> |
|---|----------------|-----------------|---------------------|---------------------|----------------------------------|-----------------|
| (10) | | | | | | |
| <i>Syniella angusta</i> Kanev | 353 | 543 | Upper Permian | Northern Urals | | |
| <i>Pseudobakewellia kolodae</i> Kanev | 366 | 10 | Lower Permian | Subpolar Urals | | |
| <i>Pseudobakewellia</i> <i>muromzevae</i> Kanev | 366 | 8 | Lower Permian | Subpolar Urals | | |
| <i>Concinnella vertnajensis</i> Kanev | 352 | 82 | Upper Permian | Preural Foredeep | | |
| <i>Mrassiella komiensis</i> Kanev | 354 | 37 | Upper Permian | Preural Foredeep | | |
| <i>Abiella chejagica</i> Kanev | 362 | 24 | Upper Permian | Pechora basin | | |
| <i>Anthraconauta (?) probus</i> Kanev | 362 | 46 | Upper Permian | Preural Foredeep | | |
| <i>Anthraconauta nativa</i> Kanev | 362 | 37 | Quarter | Pechora basin | | |
| <i>Anthraconauta uralica</i> Kanev | 356 | 425 | Upper Permian | Russian Plate | | |
| <i>Concinnella insueta</i> Kanev | 362 | 27 | Upper Permian | Pechora basin | | |