

Holotypes of fossil species

<u>Taxon</u>	<u>No coll.</u>	<u>Specimen</u>	<u>Stratigraphy</u>	<u>Location</u>	<u>Georeference</u>	<u>Bibliography</u>	<u>3D-model</u>
<u>Quarter (7)</u>							
<i>Anthraconauta nativa</i> Kanev	362	37	Quarter	Pechora basin			
<i>Loboplasma magnifica</i> Tsyganko	604	104	Quarter	Pai-Khoi			
<i>Armilla uralica</i> <i>dorsicostula</i> Sobolev	336	13-6	Quarter	Subpolar Urals			
<i>Cistula lorenziana</i> <i>var.puncta.ta</i> Losseva	192	8	Quarter	Russian Plate			
<i>Navicula insolata</i> Losseva	192	7	Quarter	Russian Plate			
<i>Raphoneis delicatus</i> (Loss.) Losseva	192	2	Quarter	Russian Plate			
<i>Pseudopodosira</i> <i>septentrionalis</i> Losseva	180	3	Quarter	Russian Plate			
<u>Neogene (3)</u>							
<i>Cyclottella notata</i> Losseva	185	18	Neogene	Russian Plate			
<i>Cyclotella omarensis</i> (Kuptz.) Loss. et Makar.	185	4	Neogene	Russian Plate			
<i>Achnanthes</i> <i>pseudoexigua</i> Losseva	185	3	Neogene	Russian Plate			