

# Holotypes of fossil species

<u>Taxon</u>	<u>№ coll.</u>	<u>Specimen</u>	<u>Stratigraphy</u>	<u>Location</u>	<u>Georeference</u>	<u>Bibliography</u>	<u>3D-model</u>
(10.)							
<i>Syniella angusta</i> Kanev	353	543	Upper Permian	Northern Urals			
<i>Pseudobakewellia kolodae</i> Kanev	366	10	Lower Permian	Subpolar Urals			
<i>Pseudobakewellia muromzevae</i> Kanev	366	8	Lower Permian	Subpolar Urals			
<i>Concinnella vertnajensis</i> Kanev	352	82	Upper Permian	Preural Foredeep			
<i>Mrassiella komiensis</i> Kanev	354	37	Upper Permian	Preural Foredeep			
<i>Abiella chejagica</i> Kanev	362	24	Upper Permian	Pechora basin			
<i>Anthraconauta (?) probus</i> Kanev	362	46	Upper Permian	Preural Foredeep			
<i>Anthraconauta nativa</i> Kanev	362	37	Quarter	Pechora basin			
<i>Anthraconauta uralica</i> Kanev	356	425	Upper Permian	Russian Plate			
<i>Concinnella insueta</i> Kanev	362	27	Upper Permian	Pechora basin			